



ELSEVIER

Comparative Biochemistry and Physiology Part B 128 (2001) 377–378

**CBP**

www.elsevier.com/locate/cbpb

## Erratum

# Erratum to “The effects of 5-HT on sensory, central and motor neurons driving the abdominal superficial flexor muscles in the crayfish”

[*Comp. Biochem. Physiol. B.* 127 (2000) 533–550]<sup>☆</sup>

J.R. Strawn<sup>a</sup>, W.S. Neckameyer<sup>b</sup>, R.L. Cooper<sup>a,\*</sup>

<sup>a</sup>*101 Thomas Hunt Morgan School of Biological Sciences, University of Kentucky,  
Lexington, KY 40506-0225, USA*

<sup>b</sup>*Department of Pharmacology and Physiological Sciences, St Louis University, St Louis,  
MO 63104, USA*

---

The publisher regrets that part B of Figure 6 was omitted from the original article and apologises for any confusion this may have caused. It is now reproduced in full on the following page:

---

<sup>☆</sup> PII of the original article: S0305-0491(00)00287-X

\* Corresponding author. Tel: +1-859-2575950; fax: +1-859-2571717.

*E-mail address:* rlcoop1@pop.uky.edu (R.L. Cooper).




Fig. 6. (A), Recording from primary sensory afferents contained within the second root of the 2nd abdominal segment. Extracellular recording illustrates the activity before cuticular stimulation, but note the large increase in the number of different size units activated during the stimulation. (B), Plots of the mean activity before and during cuticular stimulation in six preparations while bathed in saline or 5-HT (100 nM) indicates that 5-HT enhances primary sensory activity when the neurons are stimulated. This is best observed in the overall frequency of activity during the stimulation in saline and 5-HT for the same preparations.